

DIPHTHERIA TETANUS & PERTUSSIS VACCINES

WHAT YOU NEED TO KNOW

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis.

1 Why get vaccinated?

Diphtheria, tetanus, and pertussis are serious diseases caused by bacteria. Diphtheria and pertussis are spread from person to person. Tetanus enters the body through cuts or wounds.

DIPHTHERIA causes a thick covering in the back of the throat.

- It can lead to breathing problems, paralysis, heart failure, and even death.

TETANUS (Lockjaw) causes painful tightening of the muscles, usually all over the body.

- It can lead to "locking" of the jaw so the victim cannot open his mouth or swallow. Tetanus leads to death in up to 2 out of 10 cases.

PERTUSSIS (Whooping Cough) causes coughing spells so bad that it is hard for infants to eat, drink, or breathe. These spells can last for weeks.

- It can lead to pneumonia, seizures (jerking and staring spells), brain damage, and death.

Diphtheria, tetanus, and pertussis vaccine (DTaP) can help prevent these diseases. Most children who are vaccinated with DTaP will be protected throughout childhood. Many more children would get these diseases if we stopped vaccinating.

DTaP is a safer version of an older vaccine called DTP. DTP is no longer used in the United States.

2 Who should get DTaP vaccine and when?

Children should get 5 doses of DTaP vaccine, one dose at each of the following ages:

- ✓ 2 months ✓ 4 months ✓ 6 months
- ✓ 15-18 months ✓ 4-6 years

DTaP may be given at the same time as other vaccines.

3 Some children should not get DTaP vaccine or should wait

- Children with minor illnesses, such as a cold, may be vaccinated. But children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.
- Any child who had a life-threatening allergic reaction after a dose of DTaP should not get another dose.
- Any child who suffered a brain or nervous system disease within 7 days after a dose of DTaP should not get another dose.
- Talk with your doctor if your child:
 - had a seizure or collapsed after a dose of DTaP,
 - cried non-stop for 3 hours or more after a dose of DTaP,
 - had a fever over 105°F after a dose of DTaP.

Ask your health care provider for more information. Some of these children should not get another dose of pertussis vaccine, but may get a vaccine without pertussis, called **DT**.

4 Older children and adults

DTaP is not licensed for adolescents, adults, or children 7 years of age and older.

But older people still need protection. A vaccine called **Tdap** is similar to DTaP. A single dose of Tdap is recommended for people 11 through 64 years of age. Another vaccine, called **Td**, protects against tetanus and diphtheria, but not pertussis. It is recommended every 10 years. There are separate Vaccine Information Statements for these vaccines.

Diphtheria/Tetanus/Pertussis

5/17/2007

Haemophilus Influenzae Type b (Hib) Vaccine

WHAT YOU NEED TO KNOW


1 What is Hib disease?

Haemophilus influenzae type b (Hib) disease is a serious disease caused by a bacteria. It usually strikes children under 5 years old.

Your child can get Hib disease by being around other children or adults who may have the bacteria and not know it. The germs spread from person to person. If the germs stay in the child's nose and throat, the child probably will not get sick. But sometimes the germs spread into the lungs or the bloodstream, and then Hib can cause serious problems.

Before Hib vaccine, Hib disease was the leading cause of bacterial meningitis among children under 5 years old in the United States. Meningitis is an infection of the brain and spinal cord coverings, which can lead to lasting brain damage and deafness. Hib disease can also cause:

- pneumonia
- severe swelling in the throat, making it hard to breathe
- infections of the blood, joints, bones, and covering of the heart
- death



Before Hib vaccine, about 20,000 children in the United States under 5 years old got severe Hib disease each year and nearly 1,000 people died.

Hib vaccine can prevent Hib disease.

Many more children would get Hib disease if we stopped vaccinating.

2 Who should get Hib vaccine and when?

Children should get Hib vaccine at:

- ✓ 2 months of age
- ✓ 4 months of age
- ✓ 6 months of age*
- ✓ 12-15 months of age

* Depending on what brand of Hib vaccine is used, your child might not need the dose at 6 months of age. Your doctor or nurse will tell you if this dose is needed.

If you miss a dose or get behind schedule, get the next dose as soon as you can. There is no need to start over.

Hib vaccine may be given at the same time as other vaccines.

Older Children and Adults

Children over 5 years old usually do not need Hib vaccine. But some older children or adults with special health conditions should get it. These conditions include sickle cell disease, HIV/AIDS, removal of the spleen, bone marrow transplant, or cancer treatment with drugs. Ask your doctor or nurse for details.

3 Some people should not get Hib vaccine or should wait

- People who have ever had a life-threatening allergic reaction to a previous dose of Hib vaccine should not get another dose.
- Children less than 6 weeks of age should not get Hib vaccine.
- People who are moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting Hib vaccine.

Ask your doctor or nurse for more information.

POLIO VACCINE

WHAT YOU NEED TO KNOW

1 What is polio?

Polio is a disease caused by a virus. It enters a child's (or adult's) body through the mouth. Sometimes it does not cause serious illness. But sometimes it causes *paralysis* (can't move arm or leg). It can kill people who get it, usually by paralyzing the muscles that help them breathe.

Polio used to be very common in the United States. It paralyzed and killed thousands of people a year before we had a vaccine for it.

2 Why get vaccinated?

Inactivated Polio Vaccine (IPV) can prevent polio.

History: A 1916 polio epidemic in the United States killed 6,000 people and paralyzed 27,000 more. In the early 1950's there were more than 20,000 cases of polio each year. **Polio vaccination was begun in 1955.** By 1960 the number of cases had dropped to about 3,000, and by 1979 there were only about 10. The success of polio vaccination in the U.S. and other countries sparked a world-wide effort to eliminate polio.

Today: No wild polio has been reported in the United States for over 20 years. But the disease is still common in some parts of the world. It would only take one case of polio from another country to bring the disease back if we were not protected by vaccine. If the effort to eliminate the disease from the world is successful, some day we won't need polio vaccine. Until then, we need to keep getting our children vaccinated.

Oral Polio Vaccine: No longer recommended

There are two kinds of polio vaccine: **IPV**, which is the shot recommended in the United States today, and a live, oral polio vaccine (**OPV**), which is drops that are swallowed.

Until recently OPV was recommended for most children in the United States. OPV helped us rid the country of polio, and it is still used in many parts of the world.

Both vaccines give immunity to polio, but OPV is better at keeping the disease from spreading to other people. However, for a few people (about one in 2.4 million), OPV actually causes polio. Since the risk of getting polio in the United States is now extremely low, experts believe that using oral polio vaccine is no longer worth the slight risk, except in limited circumstances which your doctor can describe. The polio shot (IPV) does not cause polio. **If you or your child will be getting OPV, ask for a copy of the OPV supplemental Vaccine Information Statement.**

3 Who should get polio vaccine and when?

IPV is a shot, given in the leg or arm, depending on age. Polio vaccine may be given at the same time as other vaccines.

Children

Most people should get polio vaccine when they are children. Children get 4 doses of IPV, at these ages:

- ✓ A dose at 2 months ✓ A dose at 6-18 months
- ✓ A dose at 4 months ✓ A booster dose at 4-6 years

Adults

Most adults do not need polio vaccine because they were already vaccinated as children. But three groups of adults are at higher risk and *should* consider polio vaccination:

- (1) people traveling to areas of the world where polio is common,
- (2) laboratory workers who might handle polio virus, and
- (3) health care workers treating patients who could have polio.

Adults in these three groups who **have never been vaccinated against polio** should get 3 doses of IPV:

- ✓ The first dose at any time,
- ✓ The second dose 1 to 2 months later,
- ✓ The third dose 6 to 12 months after the second.

Adults in these three groups who **have had 1 or 2 doses** of polio vaccine in the past should get the remaining 1 or 2 doses. It doesn't matter how long it has been since the earlier dose(s).

Adults in these three groups who **have had 3 or more doses** of polio vaccine (either IPV or OPV) in the past may get a booster dose of IPV.

Ask your health care provider for more information.

4

Some people should not get IPV or should wait.

These people should not get IPV:

- Anyone who has ever had a life-threatening allergic reaction to the antibiotics **neomycin**, **streptomycin** or **polymyxin B** should not get the polio shot.
- Anyone who has a severe allergic reaction to a polio shot should not get another one.

These people should wait:

- Anyone who is moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting polio vaccine. People with minor illnesses, such as a cold, *may* be vaccinated.

Ask your health care provider for more information.

5

What are the risks from IPV?

Some people who get IPV get a sore spot where the shot was given. The vaccine used today has never been known to cause any serious problems, and most people don't have any problems at all with it.

However, a vaccine, like any medicine, could cause serious problems, such as a severe allergic reaction. *The risk of a polio shot causing serious harm, or death, is extremely small.*

6

What if there is a serious reaction?

What should I look for?

Look for any unusual condition, such as a serious allergic reaction, high fever, or unusual behavior.

If a serious allergic reaction occurred, it would happen within a few minutes to a few hours after the shot. Signs of a serious allergic reaction can include difficulty breathing, weakness, hoarseness or wheezing, a fast heart beat, hives, dizziness, paleness, or swelling of the throat.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.

- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS website at www.vaers.org, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

Reporting reactions helps experts learn about possible problems with vaccines.

7

The National Vaccine Injury Compensation Program

In the rare event that you or your child has a serious reaction to a vaccine, there is a federal program that can help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit the program's website at <http://www.hrsa.gov/osp/vicp>

8

How can I learn more?

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department's immunization program.
- Contact the Centers for Disease Control and Prevention (CDC):
-Call **1-800-232-4636 (1-800-CDC-INFO)**
-Visit the National Immunization Program's website at <http://www.cdc.gov/nip>



U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES
Centers for Disease Control and Prevention
National Immunization Program

Vaccine Information Statement

Polio (1/1/2000)

42 U.S.C. § 300aa-26